

Differential inhibition of photosynthesis under drought stress in *Flaveria* species with different degrees of development of the C₄ syndrome

M.C. DIAS and W. BRÜGGEMANN*

Department of Botany, J.W. Goethe University, POB 111932, D-60054 Frankfurt am Main, Germany

Abstract

The effect of drought stress (DS) on photosynthesis and photosynthesis-related enzyme activities was investigated in *F. pringlei* (C₃), *F. floridana* (C₃-C₄), *F. brownii* (C₄-like), and *F. trinervia* (C₄) species. Stomatal closure was observed in all species, probably being the main cause for the decline in photosynthesis in the C₃ species under ambient conditions. *In vitro* ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBPCO) and stromal fructose 1,6-bisphosphatase (sFBP) activities were sufficient to interpret the net photosynthetic rates (P_N), but, from the decreases in P_N values under high CO₂ ($C_a = 700 \mu\text{mol mol}^{-1}$) it is concluded that a decrease in the *in vivo* rate of the RuBPCO reaction may be an additional limiting factor under DS in the C₃ species. The observed decline in the photosynthesis capacity of the C₃-C₄ species is suggested to be associated both to *in vivo* decreases of RuBPCO activity and of the RuBP regeneration rate. The decline of the maximum P_N observed in the C₄-like species under DS was probably attributed to a decrease in maximum RuBPCO activity and/or to decrease of enzyme substrate (RuBP or PEP) regeneration rates. In the C₄ species, the decline of both *in vivo* photosynthesis and photosynthetic capacity could be due to *in vivo* inhibition of the phosphoenolpyruvate carboxylase (PEPC) by a twofold increase of the malate concentration observed in mesophyll cell extracts from DS plants.

Additional key words: bundle sheath cells; C₃/C₄ metabolism; CO₂ compensation concentration; malic enzyme; NADP-malate dehydrogenase; phosphoenolpyruvate carboxylase; pyruvate-orthophosphate dikinase; RuBP; RuBPCO; stromal fructose 1,6-bisphosphatase; water potential.

Introduction

The evolution of the C₄-metabolic pathway probably occurred initially about 70 million years ago as a consequence of the decrease of atmospheric CO₂ concentrations at the end of the Cretaceous period and beginning of the Paleocene, and C₄ species spread around 8–10 million years ago, when atmospheric CO₂ concentration reached a minimum of $200 \mu\text{mol mol}^{-1}$ (Ehleringer *et al.* 1991, Edwards *et al.* 2001). However, due to fossil fuel and the massive clearing of forests the concentration of atmospheric CO₂ is expected to increase up to $700 \mu\text{mol mol}^{-1}$ in the next century (Sage and Coleman 2001). Also, average atmosphere temperature is increasing and climate warming is expected to increase the frequency and severity of drought (Gregory *et al.* 1997). Drought stress (DS) is one main environmental factor limiting photosynthesis, growth, and yield of plants. Observed decreases of the

photosynthetic capacity under drought stress (DS) have been attributed to lower mesophyll CO₂ availability, as a consequence of stomatal closure (*e.g.* Chaves 1991, Cornic 2000) and/or to non-stomatal effects (*e.g.* Berkowitz 1998, Flexas and Medrano 2002, Lawlor 2002). While the majority of DS experiments have been conducted in C₃ plants, fewer have been reported from C₄ plants showing the involvement of both stomatal and non-stomatal reasons for the inhibition of photosynthesis in the latter (Du *et al.* 1996).

Environmental stresses such as DS or limited nutrient availability generally reduce the response of C₃ plants, but not of C₄ plants to increasing CO₂ concentration; this suggests that C₄ plants will continue to maintain their competitive advantages over C₃ plants under DS despite the present increase of the atmospheric CO₂ concentration

Received 2 January 2006, accepted 16 June 2006.

*Corresponding author; fax: +49-69-79824822, e-mail: w.brueggemann@bio.uni-frankfurt.de

Abbreviations: BS – bundle sheath cells; C – control; C_a – external CO₂ concentration; C_i – intercellular CO₂ concentration; DS – drought stress; FM – fresh mass; g_s – stomatal conductance; HB – homogenization buffer; MDH – malate dehydrogenase; ME – malic enzyme; P_N – net photosynthetic rate; PEPC – phosphoenolpyruvate carboxylase; PDK – pyruvate-orthophosphate dikinase; RuBP – ribulose-1,5-bisphosphate; RuBPCO – ribulose-1,5-bisphosphate carboxylase/oxygenase; sFBP – stromal fructose 1,6-bisphosphatase.

(Wand *et al.* 1999, Ward *et al.* 1999). The genus *Flaveria* (Asteraceae) contains not only C₃ and C₄ species but also various intermediates (C₃-C₄ and C₄-like species) which represent stages in the evolutionary transition from C₃ to C₄ photosynthesis (Monson and Moore 1989). We tested the hypothesis that—given comparable developmental stages, leaf properties, and a similar genetic back-

ground—the gradual development of the C₄ syndrome within a genus is beneficial for drought tolerance of photosynthesis even under nowadays or increasing CO₂ concentrations. Additionally, we compared various possible enzymatic steps for their possible role in the non-stomatal limitation of photosynthesis in the different photosynthetic pathways.

Materials and methods

Plants: All genotypes originally were kindly provided by Prof. Westhoff, University of Düsseldorf, Germany and propagated and cultivated in glasshouses of the Botanical Garden of the University of Frankfurt, Germany. Plants of *F. trinervia* (NADP-ME C₄) were grown from seeds and *F. pringlei* (C₃), *F. floridana* (C₃-C₄), and *F. brownii* (C₄-like) from cuttings in a mixture of 50 % sand and 50 % peat. After 2–3 weeks, when the cuttings presented roots, the plants were transferred to a climate chamber into plastic trays (11×11×12 cm³) containing a mixture of 1 kg(dry mass) soil (25 % sand, 25 % organic matter, 50 % peat), of which sufficient material was prepared in advance for all the experiments to ensure constant drought conditions throughout the study. The plants were grown in a climate chamber at 23 °C, 40–60 % relative humidity, 400 µmol(CO₂) mol⁻¹, 14/10 h day/night rhythm with 400 µmol(photon) m⁻² s⁻¹, provided by Osram 1000 W lamps, and received water daily. 3–4-week-old plants were exposed to DS by receiving only so much water every evening to ensure 30 % field capacity water content overnight. After 3–4 d, this resulted in estimated soil water potentials between –0.25 and –1.80 MPa during the DS treatment, which persisted for three further days.

Determination of plant water potentials: Water potentials on abscised stems was measured with a SKPM 1400 pressure chamber (SKYE Instruments, Powys, Wales, UK) according to Scholander (1965).

Gas exchange measurements: *In situ* net photosynthetic rate (P_N), stomatal conductance (g_s), and intercellular CO₂ concentration (C_i) under growth CO₂ conditions and under experimentally varied CO₂ concentrations were determined with a LI-6200 infrared gas analyzer (LiCor, Lincoln, NE, USA) at 1 000 (*F. pringlei*) or 2 000 µmol(photon) m⁻² s⁻¹ (*F. floridana*, *F. brownii*, and *F. trinervia*) “white light” either under growth chamber conditions [400 µmol(CO₂) mol⁻¹] or under experimentally varied CO₂ concentrations (Fig. 4).

Enzyme activities: Immediately after measuring the individual P_N of the youngest fully developed leaf during 15 min of irradiation [1 000 (*F. pringlei*) or 2 000 µmol (photon) m⁻² s⁻¹ (other species) for full activation of all light-regulated enzymes], enzyme activities were determined according to Du *et al.* (1996) in the same leaf

section. Two leaf discs of 11 mm diameter for *F. trinervia*, *F. pringlei*, and *F. floridana* and four leaf discs of 6 mm diameter for *F. brownii* were punched from the measured leaf section and homogenized at 0 °C with 2 cm³ homogenization buffer (HB), 20 mg Polyclar AT (SERVA, Heidelberg, Germany) and some sand with mortar and pestle. HB consisted of 50 mM Tris/HCl, pH 7.9, 8 mM MgCl₂, 5 mM Na-pyruvate, 1 mM EDTA, 2 mM K₂HPO₄, 20 mM dithiothreitol, and 0.3 % (m/v) bovine serum albumin. The homogenate was centrifuged in Eppendorf cups at 9 000×g. The supernatant was immediately used for the enzyme assays, which were performed subsequently on each homogenization before a new sample was produced from another leaf. Samples for the measurements of malate dehydrogenase (MDH), malic enzyme (ME), and phosphoenolpyruvate carboxylase (PEPC) were kept on ice until measurement. Samples for ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBPCO) determination were incubated in 20 mM MgCl₂, 10 mM NaHCO₃ for 20 min on ice prior to measurement for full activation (Lilley and Walker 1974). In the C₄ species and in both intermediate species, fully activated RuBPCO activities in the extracts usually did not match P_N . However, the same RuBPCO assay produced sufficient activities in the C₃ species. All attempts to increase the measured RuBPCO activities by variation of the homogenization procedure (grinding under liquid N₂) or increasing antioxidant or phenol-binding component concentrations failed. Since ME activities were in excess of P_N and a microscopic analysis revealed no intact bundle sheath strands in the preparations, insufficient homogenization of bundle sheath cells was excluded as a likely reason for the low RuBPCO rates. Samples for pyruvate-orthophosphate dikinase (PPDK) determination were immediately used at room temperature, since PPDK became inactivated on ice. In the case of stromal fructose 1,6-bisphosphatase (sFBP), new extracts were prepared from one leaf disc of 11 mm diameter for *F. trinervia*, *F. pringlei*, and *F. floridana* and two leaf discs of 6 mm diameter for *F. brownii* and enzyme activities were measured according to Brüggemann *et al.* (1994).

Electrophoresis: For electrophoretic and Western blot analyses of leaf extracts according to standard procedures (Laemmli 1970, Beyel and Brüggemann 2005), preparations in HB without bovine serum albumin were

performed and aliquots equivalent to defined leaf areas were run on 15 % PAA gels. Proteins were identified both by Coomassie Brilliant Blue staining and by electroblotting and incubation with anti-RuBPCO large subunit (*Secale cereale*) antiserum (Dr. Schmidt, University of Frankfurt, Germany, 1 : 5 000 dilution) or anti-PEPC (*Sorghum bicolor*) antiserum (Prof. Westhoff, University of Düsseldorf, Germany, 1 : 5 000 dilution).

Mesophyll and bundle sheath cells' separation and malate determinations in *F. trinervia*: Half of a leaf was excised and irradiated for 15 min with 2 000 μmol (photon) $\text{m}^{-2} \text{s}^{-1}$ "white light" in a moist chamber and immediately ground carefully at 0 °C with 2 cm^3 of 7 % HClO_4 and sand in a mortar and pestle to avoid bundle sheath disruption. The homogenate was then filtered through 80 μm aperture nylon net. The filtrate was neutralized with 5 M K_2CO_3 on ice and centrifuged for 5 min at 9 000 $\times g$ in an Eppendorf cup. The supernatant was termed mesophyll extract. The material remaining on the net was further ground under liquid nitrogen to ensure complete disruption of the bundle sheath cells and then

extracted with 7 % HClO_4 , neutralized with 5 M K_2CO_3 , and insoluble material pelleted by centrifugation (see above) to yield bundle sheath extract. Bundle sheath and mesophyll extracts were kept on ice until being used for the determination of malate according to Lowry and Passoneau (1972).

Microscopy analysis showed that the tissue remaining in the net contained intact bundle sheath cells and the filtrate was absent of them. Characteristic enzymes from the two cell-types (PEPC for the mesophyll cells and ME for bundle sheath cells) were determined from parallel preparations in HB instead of 7 % HClO_4 (Table 3). Enzyme activities showed a good separation by this procedure. Assuming that the filtrate mainly contained mesophyll cell fragments and the remainder in the net contained mainly intact bundle sheath cells, the fresh mass of the leaf sample used for the separation procedure was multiplied by the percentage of mesophyll (49.6 %) or bundle sheath cells (14.3 %) in a leaf of *F. trinervia*, determined from microscopy analysis, for the estimates of malate concentrations or enzyme activities for the two types of tissue.

Results

Gas exchange rates under ambient conditions: Of the four *Flaveria* species, the C_4 and the C_3 - C_4 intermediate species revealed the highest P_N under growth conditions [control (C) plants: 30–40 $\mu\text{mol m}^{-2} \text{s}^{-1}$]. Upon exposure to DS, P_N decreased with decreasing leaf water potential in all species. Drought-stressed *F. trinervia* plants (C_4) achieved the lowest plant water potentials (down to –1.9 MPa), but P_N values were above 6.5 $\mu\text{mol m}^{-2} \text{s}^{-1}$ in all cases (Fig. 1D). During DS, the lowest average P_N values were obtained in the C_3 species *F. pringlei* ($4.9 \pm 2.8 \mu\text{mol m}^{-2} \text{s}^{-1}$) (Fig. 1A).

g_s decreased in all four *Flaveria* species under DS (Fig. 2). In *F. trinervia* and *F. brownii*, the relative decrease was small due to low initial g_s . In DS plants, *F. pringlei* and *F. trinervia* presented, on average, the lowest g_s (0.030 ± 0.015 and $0.052 \pm 0.018 \text{ mol m}^{-2} \text{s}^{-1}$, respectively) (Fig. 2A,D).

The *LiCor 6200* provides an internal algorithm programme for the calculation of C_i values according to the equations developed by Caemmerer and Farquhar (1981). While these equations yield reliable values under conditions, when cuticular resistance is negligible (*i.e.* stomata are open), they tend to overestimate C_i at closed stomata, since they assume the same diffusion resistances for H_2O and CO_2 (Boyer *et al.* 1997). Nevertheless, the calculated data are summarized in Table 1.

P_N/C_i response curves: The photosynthetic response to an experimentally induced decrease of C_a was measured in C plants in the four species. While *F. pringlei* (Fig. 3A) revealed a typical RuBPCO-controlled pattern with a CO_2 compensation concentration around 50–100 $\mu\text{mol mol}^{-1}$,

Table 1. C_i values [$\mu\text{mol}(\text{CO}_2) \text{mol}^{-1}$] calculated from gas exchange data in leaves of control and drought stressed plants of *Flaveria*. Means \pm SD.

	Control	Stressed
<i>F. pringlei</i>	295 \pm 13	145 \pm 18
<i>F. floridana</i>	294 \pm 20	229 \pm 42
<i>F. brownii</i>	252 \pm 22	219 \pm 19
<i>F. trinervia</i>	256 \pm 39	224 \pm 41

F. trinervia clearly showed PEPC-controlled kinetics (Fig. 3D). In *F. pringlei* (Fig. 3B) and *F. brownii* (Fig. 3C), lower CO_2 compensation concentrations ($<50 \mu\text{mol mol}^{-1}$) were obtained with shapes of the P_N/C_i curves resembling *F. pringlei*, except for a steeper initial slope in *F. brownii*.

At high C_a (700 $\mu\text{mol mol}^{-1}$), average P_N declined under DS from 27–30 to 5–10 (*F. pringlei*), 40 to 10–20 (*F. floridana*), 25–30 to 15–20 (*F. brownii*), and 35 to 12–18 (*F. trinervia*) $\mu\text{mol}(\text{CO}_2) \text{m}^{-2} \text{s}^{-1}$, respectively.

In vitro activities of key enzymes of the Calvin cycle and electrophoretic studies: Two putative bottleneck enzymes from the Calvin cycle, RuBPCO and sFBP, were measured in all the *Flaveria* species. In the C_3 species *F. pringlei*, a decrease of about 64 % of the RuBPCO activity was observed in DS plants, but the values obtained were always above the P_N . In *F. brownii*, RuBPCO activity in DS plants was lower than in C plants, and the activities were always below the P_N . RuBPCO activities in the other two species showed




Fig. 1. Response of net photosynthetic rate (P_N) to decreasing plant water potential in control (●) and drought stressed (○) plants of *F. pringlei* (A), *F. floridana* (B), *F. brownii* (C), and *F. trinervia* (D).




Fig. 2. Response of stomatal conductance (g_s) to decreasing plant water potential in control (●) and drought stressed (○) plants of *F. pringlei* (A), *F. floridana* (B), *F. brownii* (C), and *F. trinervia* (D).

similar values under both C and DS conditions. However, these activities were insufficient to explain P_N in C plants, but exceeded P_N of DS plants (Table 2).

In accordance with the change in *in-vitro* enzyme activity, Coomassie-stained gels and Western blots of leaf extracts revealed visible decreases of the amount of RuBPCO large subunit protein only in *F. pringlei* and *F. brownii*, but not in the other two species (data not

shown).


Stromal FBP activities in both intermediates were only slightly affected by water stress. However, in *F. pringlei* and *F. trinervia* decreases of almost 50 % in the activities under DS were observed. According to the stoichiometry of the Calvin cycle (1 FBP to 3 CO_2), *in vitro* sFBP activities always exceeded the P_N in all four species (Table 2).

Table 2. RuBPCO and sFBP activities and net photosynthetic rates, P_N [$\mu\text{mol m}^{-2} \text{s}^{-1}$] in *F. pringlei*, *F. floridana*, *F. brownii*, and *F. trinervia* under control (C) and drought stress (DS) conditions. Means \pm SD.

	P_N		RuBPCO		$3 \times \text{sFBP}$	
	C	DS	C	DS	C	DS
<i>F. pringlei</i>	24.4 \pm 1.0	5.0 \pm 2.8	38.1 \pm 4.6	13.5 \pm 3.8	69.5 \pm 8.0	38.8 \pm 10.1
<i>F. floridana</i>	32.2 \pm 2.7	10.3 \pm 4.2	22.8 \pm 6.3	19.1 \pm 6.0	54.8 \pm 10.1	45.1 \pm 11.3
<i>F. brownii</i>	21.4 \pm 1.5	8.0 \pm 2.4	13.5 \pm 1.9	5.7 \pm 1.7	79.8 \pm 14.6	70.3 \pm 11.2
<i>F. trinervia</i>	36.0 \pm 1.6	10.5 \pm 2.4	19.4 \pm 4.6	19.9 \pm 4.5	81.9 \pm 11.0	43.3 \pm 6.1

Table 3. C_4 -enzyme activities and photosynthetic rates [$\mu\text{mol m}^{-2} \text{s}^{-1}$] in *F. floridana* and *F. brownii* under control (C) and drought stress (DS) conditions. Means \pm SD.

	<i>F. floridana</i>		<i>F. brownii</i>	
	C	DS	C	DS
P_N	32.2 \pm 2.7	10.3 \pm 4.2	21.4 \pm 1.5	8.0 \pm 2.4
PEPC	15.6 \pm 3.3	15.3 \pm 2.1	37.3 \pm 6.7	24.4 \pm 6.2
NADP-MDH	22.3 \pm 4.0	17.0 \pm 4.3	24.7 \pm 2.8	21.1 \pm 0.3
NADP-ME	7.4 \pm 1.5	6.1 \pm 1.2	49.7 \pm 11.7	36.5 \pm 11.4
PPDK	6.2 \pm 1.0	5.0 \pm 1.3	15.0 \pm 3.3	12.0 \pm 2.6

Fig. 3. Response of net photosynthetic rate (P_N) to intercellular CO_2 concentration (C_i) in control plants of *F. pringlei* (A), *F. floridana* (B), *F. brownii* (C), and *F. trinervia* (D).

C_4 -enzyme activities: In the C_4 plant *F. trinervia*, *in vitro* PEPC, MDH, and ME activities in both C and DS plants and PPDK activities in DS plants were always higher than the P_N (Fig. 4). However, PPDK activities in C plants were very similar to P_N . All four enzymes on average showed lower activities in DS plants than in C plants. Electrophoretic analysis of PEPC protein content revealed a slight decrease in one out of two independent preparations (data not shown).

Since both intermediate species are able to fix CO_2 at least partially through the C_4 pathway (Monson *et al.* 1986, Chastain and Chollet 1989), C_4 enzyme activities were also measured in these species. *F. brownii* always

presented higher enzyme activities than the intermediate *F. floridana*, but lower ones than the C_4 species, *F. trinervia* (Table 3, Fig. 4). The average enzyme activities measured in C and DS plants of *F. brownii* were higher than 65 % of the P_N . In *F. floridana*, the activities of the C_4 cycle enzymes measured in DS plants were at least 30 % of the P_N while in C plants only PEPC and MDH were above this percentage. In both species, only a small decrease in the C_4 enzyme activities was observed in plants under DS (Table 3). In Western blots stained with anti-PEPC antiserum, a slight decrease of PEPC protein content in leaf extracts from DS *F. brownii* was visible (data not shown).




Fig. 4. C_4 -enzyme activities compared to net photosynthetic rate (P_N) in control (●) and drought stressed (○) plants of *F. trinervia*. The solid line represents a 1 : 1 relationship between enzyme activity and photon-saturated P_N , both on leaf area basis.

Table 4. Enzyme activities [$\mu\text{mol kg}^{-1}(\text{FM}) \text{s}^{-1}$] and malate concentrations [$\text{mmol kg}^{-1}(\text{FM})$] in mesophyll and bundle sheath extracts from *F. trinervia* under control (C) and drought stress (DS) conditions. Means \pm SD.

	C		DS	
	Mesophyll	Bundle sheath	Mesophyll	Bundle sheath
PEPC	8.5 ± 1.4	4.7 ± 0.9	5.2 ± 0.9	2.6 ± 0.2
NADP-ME	1.2 ± 0.2	5.0 ± 0.3	2.3 ± 0.5	5.2 ± 0.4
Malate	35.9 ± 9.2	32.9 ± 7.6	72.3 ± 15.8	35.9 ± 9.2

Malate contents in mesophyll and bundle sheath cells from *F. trinervia*: Malate is a known inhibitor of PEPC (Wedding *et al.* 1990). In order to study whether a putative malate feedback inhibition may occur under DS, as postulated for *Sorghum bicolor* (Beyel and Brüggemann

2005), mesophyll and bundle sheath cell extracts were prepared and malate contents were measured. Malate contents increased twofold in mesophyll extracts of DS plants. However, in bundle sheath extracts, malate contents in C and DS plants were similar (Table 4).

Discussion

Stomatal vs. non-stomatal effects: A decrease in P_N during water deficit can be attributed to both stomatal and non-stomatal effects. Stomatal closure is considered the first line of defence against water loss (Chaves 1991), and a parallel decrease of P_N and g_s under DS has generally been reported (Medrano *et al.* 1997, Correia *et al.* 1999, Castrillo *et al.* 2001, Maroco *et al.* 2002). Also in the four *Flaveria* species of this study, DS lead to stomatal closure (Fig. 2). As in other C_3 species (Sharkey and Seemann 1989, Vassey and Sharkey 1989, Lal *et al.* 1996, Escalona *et al.* 1999), stomatal closure strongly reduced the intercellular CO_2 availability in *F. pringlei* (Table 1). In the other three species, the *LiCor* calculation revealed no significant decrease of C_i , in accordance with previous studies on other C_4 plants (Premachandra *et al.* 1994, Prakash and Rao 1996, Yu *et al.* 2004, Beyel and Brüggemann 2005). However, the reliance of C_i calcu-

lations from CO_2 and H_2O exchange rates in DS plants is strongly under debate, not only because of the masked effects of stomatal patchiness (Terashima *et al.* 1988, Mansfield *et al.* 1990, Terashima 1992), but also because of the different diffusion resistances of the cuticle for CO_2 and H_2O (Boyer *et al.* 1997). Therefore, no unequivocal conclusions on possible stomatal limitations of photosynthesis could be drawn for the other three species, and measurements of photosynthetic enzymes were performed to identify possible non-stomatal limiting factors.

In *F. pringlei*, fully activated *in vitro* RuBPCO activity and RuBPCO protein content decreased in DS plants, although the activities still exceeded the P_N . Since the calculated decline of C_i alone would still allow for P_N higher than those observed, provided the calculations were reliable and the P_N/C_i curve for C plants (Fig. 3A)

would still be applicable to DS plants, a decline of the *in vivo* RuBPCO activity may well contribute to the limitation of photosynthesis under DS. Similarly, Kicheva *et al.* (1994) in wheat, Kanechi *et al.* (1996) in coffee, and Panković *et al.* (1999) in sunflower observed decreases of *in vitro* RuBPCO activity under DS. In principle, several factors may contribute to even further declines of RuBPCO activity *in vivo* through side effects of DS: stromal acidification (Berkowitz *et al.* 1983), increase of stromal ionic strength (Kaiser 1982), or accumulating RuBPCO oxidation through over-excitation, which can play a role under long-term chilling stress (Brüggemann *et al.* 1995). Limitation of RuBP regeneration by the decreased sFBP activity (Table 2) appears unlikely, since the latter enzyme activity still vastly exceeded the P_N .

Positive P_N in the C_3 - C_4 intermediate and the C_4 -like species were already observed at C_i below $50 \mu\text{mol mol}^{-1}$ (Fig. 4B,C). These results are attributed to reduced photorespiratory loss of CO_2 by (partially expressed) C_4 -metabolism (*F. brownii*) and by recycling the photorespired CO_2 in the bundle sheath cells in the case of C_3 - C_4 intermediate species (*F. floridana*), respectively (Monson 1989). C_3 - C_4 intermediate species are able to re-assimilate up to 70 % of the photorespired CO_2 (Hunt *et al.* 1987). In the C_4 -like intermediate species this mechanism is not so important since the reduced photorespiration rates are mainly due to the high level of development of the C_4 pathway (Cheng *et al.* 1988, Ku *et al.* 1991).

RuBPCO activities in C plants of *F. floridana* and *F. brownii* were lower than the respective P_N . In *Sorghum*, where a similar finding was observed, Western blots gave no indications of a decrease of RuBPCO protein content in DS plants (Beyel and Brüggemann 2005). According to Rogers *et al.* (2001), who studied pine needle tissue, insufficient extraction of RuBPCO before the analysis may often be the cause for low RuBPCO activities *in vitro* in leaves with strong cell walls. However, any attempt to increase *in vitro* RuBPCO activities of C plants to or beyond their photosynthetic capacity by modifying the extraction procedure (liquid N_2 and excessive grinding) failed both in *Sorghum* (Beyel and Brüggemann 2005) and in *Flaveria*.

F. floridana exhibits characteristics between the C_3 and C_4 pathways (Holaday *et al.* 1984). RuBPCO is present in mesophyll and bundle sheath cells while PEPC and NADP-ME are partially compartmentalized as in a C_4 plant (Casati *et al.* 1999). Since the intermediate species contains both C_3 and C_4 pathway activities, both enzymes, RuBPCO and PEPC, are involved in the first step of the CO_2 assimilation. However, in *F. floridana*, PEPC is only responsible for a portion of the total CO_2 fixed (15 % according to Holaday *et al.* 1984, 30–50 % according to Chastain and Chollet 1989 and to Monson *et al.* 1986). Our results exclude the C_4 enzymes as possible limiting factors, since the activities measured in C and DS plants were similar and were enough to explain above 30 % of P_N . In *F. floridana*, the P_N/C_i curve strongly

points to a RuBPCO-reaction limited situation of P_N in C plants under most CO_2 concentrations, be it by CO_2 or by RuBP regeneration (Fig. 3B). *In vitro* RuBPCO activities as well as RuBPCO protein contents were similar in C and DS plants (Table 2). While under C_a , stomatal limitation alone may well account for the observed decline in P_N , the decrease of P_N even under higher CO_2 in DS plants of *F. floridana* could be associated to an *in vivo* decrease in the RuBP regeneration cycle. Again, however, measurements of sFBP in *F. floridana* showed only a small decrease in the *in vitro* enzyme activity under DS, probably ruling this enzyme out as a limiting factor as discussed for *F. pringlei* (Table 2).

F. brownii is the most advanced C_3 - C_4 intermediate in the genus *Flaveria* in terms of development of the C_4 syndrome (Monson *et al.* 1987). However, the principal photosynthetic carboxylation enzymes are not yet completely compartmentalized between mesophyll and bundle sheath cells as in a C_4 plant. PEPC and NADP-ME enzyme are present in both mesophyll and bundle sheath cells, but RuBPCO is mainly present in the bundle sheath cells chloroplasts (Reed and Chollet 1985). In *F. brownii*, about 65–80 % of the CO_2 is fixed into C_4 acids, whereas about 20 % of CO_2 enters the C_3 cycle directly through RuBPCO (Monson *et al.* 1986, Cheng *et al.* 1988, Chastain and Chollet 1989). The analysis of the P_N/C_i curve and the activity of PEPC suggest that the kinetic of CO_2 uptake at low CO_2 (initial slope) is related to the activity of PEPC as in a C_4 plant, while for high CO_2 it represents either the *in vivo* RuBPCO activity or the *in vivo* PEP regeneration rate (Caemmerer and Furbank 1999). The decline of the maximum P_N in DS plants under elevated CO_2 was in parallel to the decrease in the maximum RuBPCO activity observed in the *in vitro* measurements. Under DS, both PEPC activity and PEPC protein content declined slightly. However, DS plants of *F. brownii* contained sufficient *in vitro* PEPC, MDH, and ME activities to exceed the P_N under elevated CO_2 concentration (Table 3), while PPDK might become limiting in addition to the effect of DS on RuBPCO. Thus, with stomata closed in DS plants, the plants may still perform the P_N allowed by (less) PEPC, but will lose the additional beneficial effect of RuBPCO at higher C_i (cf. P_N/C_i curve, Fig. 3C).

In *F. trinervia*, both P_N under ambient CO_2 and under elevated CO_2 decreased to similar values [$10\text{--}15 \mu\text{mol}(\text{CO}_2) \text{m}^{-2} \text{s}^{-1}$] when plants were exposed to DS. The decrease in maximum P_N may reflect decreases in RuBPCO reaction rate and/or in the *in vivo* PEP regeneration rate (cf. Figs. 5 and 6 in Caemmerer and Furbank 1999). However, measurements of the *in vitro* activities of RuBPCO and of PEPC and other enzymes of the C_4 -metabolism showed that all major enzymes were sufficient to explain photosynthetic activities of the DS plants (Fig. 4). Only in the case of PPDK in C plants, this enzyme may be a limiting factor for photosynthetic rates. Du *et al.* (1996) also observed a decrease of the PPDK

activity in DS sugarcane plants. In DS plants the two Calvin cycle enzymes, sFBP and RuBPCO, also revealed activities in excess of P_N . However, as in *F. floridana* and *F. brownii*, in C plants the values for RuBPCO activity were below the P_N .

Malate concentrations in *F. trinervia*: The results of the gas exchange measurements had indicated that the decline of the photosynthesis capacity in *F. trinervia* could be due to an inhibition of the CO₂-fixing enzyme PEPC *in vivo* under DS, despite sufficient PEPC activity *in vitro*. PEPC is the key enzyme of carbon assimilation in the C₄ pathway and is located in the cytoplasm of mesophyll cells. The activity of this enzyme is regulated by metabolites, changes in the pH of the cytoplasm, and also by reversible phosphorylation of a serine residue, which alters the sensitivity of the enzyme to these metabolites (Nimmo *et al.* 1986, Chollet *et al.* 1996, Vidal and Chollet 1997). While malate, the major product of the primary CO₂ fixation, is a competitive inhibitor of the PEPC activity, glucose-6-phosphate increases the affinity of PEPC for PEP by reducing the K_m , this positive effector enhances the ability of PEP to compete with malate (Vidal and Chollet 1997). Phosphorylation of PEPC in irradiated leaves markedly increases PEPC activity, reduces the inhibitory effect of malate, and increases the effect of glucose-6-phosphate (Leegood and Walker 1999). Since the observed *in vitro* PEPC activities in DS plants of *F. trinervia* vastly exceeded P_N without protein kinase and ATP in the extraction medium (*i.e.* a technique used by Sawada *et al.* 2002 to assess the phosphorylation state *in vivo*), we concluded that a putative decrease in the *in vivo* phosphorylation state of the enzyme was not responsible for the decline in P_N . Enzyme phosphorylation seems to allow C₄ PEPC to continue to fix CO₂ in the presence of high cytosolic malate concentrations (10–20 mM) in mesophyll cells in order to maintain malate diffusion to bundle sheath cells (Vidal and Chollet 1997). As observed here in *F. trinervia* and also by other authors in *Zea mays* (Leegood 1985, Stitt and Heldt 1985), malate is mainly located in mesophyll cells. However, not all malate is photosynthetically active in the cytoplasm (Hatch 1971), but a considerable amount is probably stored in the vacuole (Leegood 1985). In the literature, malate contents measured in C₄ leaves (*Zea mays*) range from 5–17 (bundle sheath cells) to 35–92 mM (mesophyll cells) (Leegood 1985, Stitt and Heldt 1985). Beyel and Brüggemann (2005) observed *ca.* 80 % increase in total leaf malate content in *Sorghum bicolor* under DS. Different results were observed by Du *et al.* (1998) in sugarcane, where malate contents dropped in leaves of DS plants. In *F. trinervia*, malate contents in the mesophyll extracts doubled from 36 in C up to 72 mmol kg⁻¹(FM) in DS plants. In bundle sheath extracts, these values were similar in C [32 mmol kg⁻¹(FM)] and DS plants [36 mmol kg⁻¹(FM)]. However, we did not observe a sig-

nificant gradient of overall malate contents between BS and mesophyll cells of C plants, nor did we observe a significant increase in overall malate contents in the BS upon DS. While, therefore, conclusions from the overall contents upon the (critical) cytoplasm contents should be made with care, the data clearly indicate a significant increase of whole cell malate content in the mesophyll as a consequence of feedback accumulation through limited sink activity under DS.

C₄-species of the genus *Flaveria* contain both C₃ and C₄ isoforms of the enzyme PEPC; the C₃ form of PEPC showing a 15-fold higher sensitivity to malate than the C₄-enzyme (Bläsing *et al.* 2002). However, when the pH decreases to 7.3, also the C₄ isoform becomes inactivated at a malate content of 9.4 mM (Bläsing *et al.* 2002). Under photosynthetic conditions, the cytosolic pH undergoes a moderate acidification by dicarbonic acid formation (Yin *et al.* 1993), and DS decreases cytosolic pH further (Berkowitz *et al.* 1983). Even if the malate content in the cytoplasm would only be 10 % of the overall concentration found in mesophyll cells of DS plants (*i.e.* 7 mM, Table 4), at pH 7.3 or lower it would be strongly inhibiting.

Concluding remarks: The results indicate differential sensitivity of photosynthesis towards DS in the four species, depending on different physiological factors. An increase of malate concentration in the mesophyll cells may be the cause for the inhibition of PEPC activity *in vivo* and photosynthesis in the C₄ species. The decline of photosynthesis in the C₄-like intermediate was most probably due to a combination of stomatal closure and partial PEPC and RuBPCO losses in DS plants, while in the C₃-C₄ intermediate species it was probably associated with a reduction of g_s and, possibly, the regenerative phase of the Calvin cycle at elevated CO₂ concentration. In the C₃ species, the decrease in photosynthesis under DS was probably related to both (complete) stomatal closure and to decreases in the rate of the RuBPCO reaction.

The positive P_N observed at very low C_i (<50 μmol mol⁻¹) in both intermediate species represents an advantage under low atmospheric CO₂ when compared to the C₃ species.

Although non-stomatal effects severely decreased P_N in DS plants of the other three species, their P_N under C_a was higher than in *F. pringlei*. As shown in Fig. 2, the dependence of CO₂ fixation on the enzymatic properties of RuBPCO forces the C₃ and C₃-C₄ species to operate at higher g_s under C conditions, leading to lower water use efficiency than in the C₄ and C₄-like species. Under DS, the evolutionary change from a RuBPCO-based to a PEPC-based CO₂ fixing system *via* the C₃-C₄ intermediate and the C₄-like type, results in a secondary advantage, *i.e.* to produce sufficient saccharides to maintain the reduced need of the sinks even with stomata closed.

Since the C₃-C₄ and C₄-like species represent evolutionary intermediates that have evolved from C₃ plants

(Monson and Moore 1989), our results indicate that even under today's high atmospheric CO₂ concentration, the further development of the C₄ pathway in these inter-

mediates will still be beneficial under water-limiting growth conditions.

References

- Berkowitz, G.A.: Water and salt stress. – In: Raghavendra, A.S. (ed.): *Photosynthesis. A Comprehensive Treatise*. Pp. 226-237. Cambridge University Press, Cambridge 1998.
- Berkowitz, G.A., Chen, C., Gibbs, M.: Stromal acidification mediates *in vivo* water stress inhibition of nonstomatal-controlled photosynthesis. – *Plant Physiol.* **72**: 1123-1126, 1983.
- Beyel, V., Brüggemann, W.: Differential inhibition of photosynthesis during pre-flowering drought stress in *Sorghum bicolor* genotypes with different senescence traits. – *Physiol. Plant.* **124**: 249-259, 2005.
- Bläsing, O., Ernst, K.P., Streubel, M., Westhooff, P., Svensson, P.: The non-photosynthetic phosphoenolpyruvate carboxylases of the C₄ dicot *Flaveria trinervia* – implications for the evolution of C₄ photosynthesis. – *Planta* **215**: 448-456, 2002.
- Boyer, J.S., Wong, S.C., Farquhar, G.D.: CO₂ and water vapor exchange across leaf cuticle (epidermis) at various water potentials. – *Plant Physiol.* **114**: 185-191, 1997.
- Brüggemann, W., Dauborn, B., Klaucke, S., Linger, P., Maas-Kantel, K., Wenner, A.: Chilling sensitivity of photosynthesis: Ecophysiological studies in two *Lycopersicon* species of different chilling tolerance. – *Acta Physiol. Plant.* **17**: 113-122, 1995.
- Brüggemann, W., Klaucke, S., Maas-Kantel, K.: Long-term chilling of young tomato plants under low light. V. Kinetic and molecular properties of two key enzymes of the Calvin cycle in *Lycopersicon esculentum* Mill. and *L. peruvianum* Mill. – *Planta* **194**: 160-168, 1994.
- Caemmerer, S. von, Farquhar, G.D.: Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. – *Planta* **153**: 376-387, 1981.
- Caemmerer, S. von, Furbank, R.T.: Modelling C₄ photosynthesis. – In: Sage, R.F., Monson, R.K. (ed.): *C₄ Plant Biology*. Pp. 173-211. Academic Press, San Diego 1999.
- Casati, P., Fresco, A., Andreo, C., Drincovich, M.F.: An intermediate form of NADP-malic enzyme from the C₃-C₄ intermediate species *Flaveria floridana*. – *Plant Sci.* **147**: 101-109, 1999.
- Castrillo, M., Fernandez, D., Calcagno, A.M., Trujillo, I., Guenni, L.: Responses of ribulose-1,5-bisphosphate carboxylase, protein content, and stomatal conductance to water deficit in maize, tomato, and bean. – *Photosynthetica* **39**: 221-226, 2001.
- Chastain, C., Chollet, R.: Interspecific variation in assimilation of ¹⁴CO₂ into C₄ acids by leaves of C₃, C₄ and C₃-C₄ intermediate *Flaveria* species near the CO₂ compensation concentration. – *Planta* **179**: 81-88, 1989.
- Chaves, M.M.: Effects of water deficits on carbon assimilation. – *J. exp. Bot.* **42**: 1-16, 1991.
- Cheng, S.-H., Moore, B.D., Edwards, G.E., Ku, M.S.B.: Photosynthesis in *Flaveria brownii*, a C₄-like species. Leaf anatomy, characteristics of CO₂ exchange, compartmentation of photosynthetic enzymes, and metabolism of ¹⁴CO₂. – *Plant Physiol.* **87**: 867-873, 1988.
- Chollet, R., Vidal, J., O'Leary, M.H.: Phosphoenolpyruvate carboxylase: A ubiquitous, highly regulated enzyme in plants. – *Annu. Rev. Plant Physiol. Plant mol. Biol.* **47**: 273-298, 1996.
- Cornic, G.: Drought stress inhibits photosynthesis by decreasing stomatal aperture – not by affecting ATP synthesis. – *Trends Plant Sci.* **5**: 187-188, 2000.
- Correia, M.J., Rodrigues, M.L., Osório, M.L., Chaves, M.M.: Effects of growth temperature on the response of lupin stomata to drought and abscisic acid. – *Aust. J. Plant Physiol.* **26**: 549-559, 1999.
- Du, Y.C., Kawamitsu, Y., Nose, A., Hiyane, S., Murayama, S., Wasano, K., Uchida, Y.: Effects of water stress on carbon exchange rate and activities of photosynthetic enzymes in leaves of sugarcane (*Saccharum* sp.). – *Aust. J. Plant Physiol.* **23**: 719-726, 1996.
- Du, Y.-C., Nose, A., Wasano, K., Uchida, Y.: Responses to water stress of enzyme activities and metabolite levels in relation to sucrose and starch synthesis, the Calvin cycle and the C₄ pathway in sugarcane (*Saccharum* sp.) leaves. – *Aust. J. Plant Physiol.* **25**: 253-260, 1998.
- Edwards, G.E., Furbank, R.T., Hatch, M.D., Osmond, C.B.: What does it take to be C₄? Lessons from the evolution of C₄ photosynthesis. – *Plant Physiol.* **125**: 46-49, 2001.
- Ehleringer, J.R., Sage, R.F., Flanagan, L.B., Pearcy, R.W.: Climate change and the evolution of C₄ photosynthesis. – *Trends Ecol. Evolut.* **6**: 95-99, 1991.
- Escalona, J.M., Flexas, J., Medrano, H.: Stomatal and non-stomatal limitations of photosynthesis under water stress in field-grown grapevines. – *Aust. J. Plant Physiol.* **26**: 421-433, 1999.
- Flexas, J., Medrano, H.: Drought-inhibition of photosynthesis in C₃ plants: Stomatal and non-stomatal limitations revisited. – *Ann. Bot.* **89**: 183-189, 2002.
- Gregory, J.M., Mitchell, J.F.B., Brady, A.J.: Summer drought in northern midlatitudes in a time-dependent CO₂ climate experiment. – *J. Climate* **10**: 662-686, 1997.
- Hatch, M.D.: The C₄-pathway of photosynthesis. Evidence for an intermediate pool of carbon dioxide and the identity of the donor C₄-dicarboxylic acid. – *Biochem. J.* **125**: 425-432, 1971.
- Holaday, A.S., Lee, K.W., Chollet, R.: C₃-C₄ intermediate species in the genus *Flaveria*: leaf anatomy, ultrastructure, and the effect of O₂ on the CO₂ compensation concentration. – *Planta* **160**: 25-32, 1984.
- Hunt, S., Smith, A.M., Woolhouse, H.W.: Evidence for a light-dependent system for reassimilation of photorespiratory CO₂, which does not include a C₄ cycle, in the C₃-C₄ intermediate species *Moricandia arvensis*. – *Planta* **171**: 227-234, 1987.
- Kaiser, W.M.: Correlation between changes in photosynthetic activity and changes in total protoplast volume in leaf tissue from hygro-, meso- and xerophytes under osmotic stress. – *Planta* **154**: 538-545, 1982.
- Kanечи, M., Uchida, N., Yasuda, T., Yamaguchi, T.: Non-stomatal inhibition associated with inactivation of Rubisco in dehydrated coffee leaves under unshaded and shaded conditions. – *Plant Cell Physiol.* **37**: 455-460, 1996.
- Kicheva, M.I., Tsonev, T.D., Popova, L.P.: Stomatal and non-stomatal limitations to photosynthesis in two wheat cultivars subjected to water stress. – *Photosynthetica* **30**: 107-116, 1994.

- Ku, M.S.B., Wu, J., Dai, Z., Scott, R.A., Chu, C., Edwards, G.E.: Photosynthetic and photorespiratory characteristics of *Flaveria* species. – *Plant Physiol.* **96**: 518-528, 1991.
- Laemmli, U.K.: Cleavage of structural proteins during the assembly of the head of bacteriophage T4. – *Nature* **227**: 680-685, 1970.
- Lal, A., Ku, M.S.B., Edwards, G.E.: Analysis of inhibition of photosynthesis due to water stress in the C₃ species *Hordeum vulgare* and *Vicia faba*: Electron transport, CO₂ fixation and carboxylation activity. – *Photosynth. Res.* **49**: 57-69, 1996.
- Lawlor, D.W.: Limitation to photosynthesis in water-stressed leaves: Stomata vs. metabolism and the role of ATP. – *Ann. Bot.* **89**: 871-885, 2002.
- Leegood, R.C.: The intercellular compartmentation of metabolites in leaves of *Zea mays* L. – *Planta* **164**: 163-171, 1985.
- Leegood, R.C., Walker, R.P.: Regulation of the C₄ pathway. – In: Sage, R.F., Monson, R.K. (ed.): *C₄ Plant Biology*. Pp. 89-131. Academic Press, San Diego 1999.
- Lilley, R.M.C., Walker, D.A.: An improved spectrophotometric assay for ribulosebiphosphate carboxylase. – *Biochim. biophys. Acta* **358**: 226-229, 1974.
- Lowry, O.H., Passoneau, J.V.: *A Flexible System of Enzymatic Analysis*. – Academic Press, London 1972.
- Mansfield, T.A., Hetherington, A.M., Atkinson, C.J.: Some current aspects of stomatal physiology. – *Annu. Rev. Plant Physiol. Plant mol. Biol.* **41**: 55-75, 1990.
- Maroco, P.M., Rodrigues, M.L., Lopes, C., Chaves, M.M.: Limitations to leaf photosynthesis in field-grown grapevine under drought – metabolic and modelling approaches. – *Funct. Plant Biol.* **29**: 451-459, 2002.
- Medrano, H., Parry, M.A.J., Socías, X., Lawlor, D.W.: Long term water stress inactivates Rubisco in subterranean clover. – *Ann. appl. Biol.* **131**: 491-501, 1997.
- Monson, R.K.: The relative contributions of reduced photorespiration, and improved water- and nitrogen-use efficiencies, to the advantages of C₃-C₄ intermediate photosynthesis in *Flaveria*. – *Oecologia* **80**: 215-221, 1989.
- Monson, R.K., Moore, B.D.: On the significance of C₃-C₄ intermediate photosynthesis to the evolution of C₄ photosynthesis. – *Plant Cell Environ.* **12**: 689-699, 1989.
- Monson, R.K., Moore, B.D., Ku, M.S.B., Edwards, G.E.: Co-function of C₃- and C₄-photosynthetic pathways in C₃, C₄ and C₃-C₄ intermediate *Flaveria* species. – *Planta* **168**: 493-502, 1986.
- Monson, R.K., Schuster, W.S., Ku, M.S.B.: Photosynthesis in *Flaveria brownii* A.M. Powell. A C₄-like C₃-C₄ intermediate. – *Plant Physiol.* **85**: 1063-1067, 1987.
- Nimmo, G.A., Nimmo, H.G., Hamilton, I.D., Fewson, C.A., Wilkins, M.B.: Purification of the phosphorylated night form and dephosphorylated day form of phosphoenolpyruvate carboxylase from *Bryophyllum fedtschenkoi*. – *Biochem. J.* **239**: 213-220, 1986.
- Panković, D., Sakač, Z., Kevrešan, S., Plesničar, M.: Acclimation to long-term water deficit in the leaves of two sunflower hybrids: photosynthesis, electron transport and carbon metabolism. – *J. exp. Bot.* **50**: 127-138, 1999.
- Prakash, K.R., Rao, V.S.: The altered activities of carbonic anhydrase, phosphoenol pyruvate-carboxylase and ribulose-bisphosphate carboxylase due to water-stress and after its relief. – *J. environ. Biol.* **1**: 39-42, 1996.
- Premachandra, G.S., Hahn, D.T., Joly, R.J.: Leaf water relations and gas exchange in two grain *Sorghum* genotypes differing in their pre- and post-flowering drought tolerance. – *J. Plant Physiol.* **143**: 96-101, 1994.
- Reed, J.E., Chollet, R.: Immunofluorescent localization of phosphoenolpyruvate carboxylase and ribulose 1,5-bisphosphate carboxylase/oxygenase proteins in leaves of C₃, C₄ and C₃-C₄ intermediate *Flaveria* species. – *Planta* **165**: 439-445, 1985.
- Rogers, A., Ellsworth, D.S., Humphries, S.W.: Possible explanation of the disparity between the *in vitro* and *in vivo* measurements of Rubisco activity: a study in loblolly pine grown in elevated pCO₂. – *J. exp. Bot.* **52**: 1555-1561, 2001.
- Sage, R.F., Coleman, J.R.: Effects of low atmospheric CO₂ on plants: more than a thing of the past. – *Trends Plant Sci.* **6**: 18-24, 2001.
- Sawada, S., Sakamoto, T., Sato, M., Kasai, M., Usuda, H.: Photosynthesis with single-rooted *Amaranthus* leaves. II. Regulation of ribulose-1,5-bisphosphate carboxylase, phosphoenolpyruvate carboxylase, NAD-malic enzyme and NAD-malate dehydrogenase and coordination between PCR and C₄ photosynthetic metabolism in response to changes in the source-sink balance. – *Plant Cell Physiol.* **43**: 1293-1301, 2002.
- Sharkey, T.D., Seemann, J.R.: Mild water stress effects on carbon-reduction-cycle intermediates, ribulose bisphosphate carboxylase activity, and spatial homogeneity of photosynthesis in intact leaves. – *Plant Physiol.* **89**: 1060-1065, 1989.
- Scholander, P.F.: Sap pressure in vascular plants. – *Science* **148**: 339-346, 1965.
- Stitt, M., Heldt, H.W.: Generation and maintenance of concentration gradients between the mesophyll and bundle sheath in maize leaves. – *Biochim. biophys. Acta* **808**: 400-414, 1985.
- Terashima, I.: Anatomy of non-uniform leaf photosynthesis. – *Photosynth. Res.* **31**: 195-212, 1992.
- Terashima, I., Wong, S.-C., Osmond, C.B., Farquhar, G.D.: Characterisation of non-uniform photosynthesis induced by abscisic acid in leaves having different mesophyll anatomies. – *Plant Cell Physiol.* **29**: 385-394, 1988.
- Vassey, T.L., Sharkey, T.D.: Mild water stress of *Phaseolus vulgaris* plants leads to reduced starch synthesis and extractable sucrose phosphate synthase activity. – *Plant Physiol.* **89**: 1066-1070, 1989.
- Vidal, J., Chollet, R.: Regulatory phosphorylation of C₄ PEP carboxylase. – *Trends Plant Sci.* **2**: 230-241, 1997.
- Wand, S.J.E., Midgley, G.F., Jones, M.H., Curtis, P.S.: Responses of wild C₄ and C₃ grass (Poaceae) species to elevated atmospheric CO₂ concentrations: a meta-analytic test of current theories and perceptions. – *Global Change Biol.* **5**: 723-741, 1999.
- Ward, J.K., Tissue, D.T., Thomas, R.B., Strain, R.B.: Comparative responses of model C₃ and C₄ plants to drought in low and elevated CO₂. – *Global Change Biol.* **5**: 857-867, 1999.
- Wedding, R.T., Black, M.K., Meyer, C.R.: Inhibition of phosphoenolpyruvate carboxylase by malate. – *Plant Physiol.* **92**: 456-461, 1990.
- Yin, Z.-H., Heber, U., Raghavendra, A.S.: Light-induced pH changes in leaves of C₄ plants. Comparison of cytosolic alkalization and vacuolar acidification with that of C₃ plants. – *Planta* **189**: 267-277, 1993.
- Yu, G., Wang, Q., Zhuang, J.: Modelling the water use efficiency of soybean and maize plants under environmental stresses: application of a synthetic model of photosynthesis-transpiration based on stomatal behavior. – *J. Plant Physiol.* **161**: 303-318, 2004.